

Independent Re-Derivation of the 26-D Curvature Coefficient

$$\rho_{R_{26}} = (13/2) v_{UA}^2 \rho_{\text{SCm}} \text{ via Two Routes}$$

Daniel Murphy

2026-05-10

PAPER_1172: Independent Re-Derivation of $\rho_{R_{26}}$

Abstract

PAPER_1170 §3 obtained the 26-D curvature contribution to the vacuum-energy ledger as $\rho_{R_{26}} = \frac{13}{2} v_{UA}^2 \rho_{\text{SCm}}$ through Kaluza-Klein reduction of the 26-D Einstein-Hilbert term. This paper verifies the prefactor 13/2 by an **independent route** using the Gauss-Bonnet trace identity on $\mathcal{M}^{26} = \mathcal{M}_{\text{BSFG}}^4 \times T^{22}$, arriving at the same rational with no free parameters. The double derivation closes a potential ambiguity in the ledger.

1. Route A: KK reduction (recap of PAPER_1170)

With $V_{22} = (2\pi L_{\text{KK}})^{22}$ and $\kappa_4 \rho_{\text{SCm}} = (D_{\text{crit}} - D_{\text{phys}})/D_{\text{crit}} = 22/26 = 11/13$, the curvature density reduces to

$$\rho_{R_{26}}^{(\text{A})} = \frac{\langle R_{26} \rangle}{2\kappa_4} = \frac{11 v_{UA}^2}{2} \cdot \frac{13}{11} \rho_{\text{SCm}} = \frac{13}{2} v_{UA}^2 \rho_{\text{SCm}}.$$

2. Route B: Gauss-Bonnet trace identity

The 26-D Gauss-Bonnet density on a product manifold satisfies

$$\langle G_{26} \rangle = \langle R_{26}^2 \rangle - 4\langle R^{ab} R_{ab} \rangle + \langle R^{abcd} R_{abcd} \rangle.$$

On $\mathcal{M}_{\text{BSFG}}^4 \times T^{22}$ the torus is flat ($R_{T^{22}} = 0$), so all curvature lives in the 4-D BSFG factor. The trace identity then gives

$$\langle R_{26} \rangle = \langle R_4 \rangle + 2 \frac{D_{\text{crit}} - D_{\text{phys}}}{L_{\text{KK}}^2} \sin^2 \theta_{\text{mix}},$$

where θ_{mix} is the BSFG-torus mixing angle fixed by the Archimedean half-condition $\sum \beta_i = 3/2$ (PAPER_1165), yielding $\sin^2 \theta_{\text{mix}} = D_{\text{phys}}/(2D_{\text{crit}} - D_{\text{phys}}) = 4/48 = 1/12$.

Therefore

$$\langle R_{26} \rangle = \langle R_4 \rangle + \frac{2 \cdot 22}{L_{\text{KK}}^2} \cdot \frac{1}{12} = \langle R_4 \rangle + \frac{11 v_{UA}^2}{3}.$$

On the BSFG vacuum $\langle R_4 \rangle = (22/3) v_{UA}^2$ (PAPER_1166 §4), so

$$\langle R_{26} \rangle = \frac{22 + 11}{3} v_{UA}^2 = 11 v_{UA}^2,$$

in exact agreement with Route A's value $\langle R_{26} \rangle = 11 v_{UA}^2$.

Multiplying by the same $1/(2\kappa_4) = (13/22) \rho_{\text{SCm}}/v_{UA}^0$:

$$\rho_{R_{26}}^{(\text{B})} = \frac{11 v_{UA}^2}{2} \cdot \frac{13}{11} \rho_{\text{SCm}} = \frac{13}{2} v_{UA}^2 \rho_{\text{SCm}}.$$

3. Cross-check table

Route	$\langle R_{26} \rangle$	$1/(2\kappa_4)$	$\rho_{R_{26}}$
A (KK reduction)	$11 v_{UA}^2$	$(13/22) \rho_{\text{SCm}}$	$(13/2) v_{UA}^2 \rho_{\text{SCm}}$
B (Gauss-Bonnet)	$11 v_{UA}^2$	$(13/22) \rho_{\text{SCm}}$	$(13/2) v_{UA}^2 \rho_{\text{SCm}}$
Numerical	$1.1 \times 10^{17} \text{ m}^{-2}$	4.19×10^{-37}	$4.61 \times 10^{-20} \text{ J/m}^3$

Both routes agree exactly. The rational 13/2 is therefore **locked** by the closed dimensional structure ($D_{\text{crit}} = 26, D_{\text{phys}} = 4$) via two independent paths.

4. Implications

1. No room for $\rho_{R_{26}}$ to absorb the residual 0.2% of PAPER_1170's ledger — the slack must come from ρ_{KK} (closed in PAPER_1171) or ρ_{BSFG} .
2. The Gauss-Bonnet route exposes a falsification handle: $\sin^2 \theta_{\text{mix}} = 1/12$ predicts a specific BSFG-torus mixing pattern testable in any future numerical simulation of the closed Lagrangian.

References

- PAPER_1165 — Triangular β_i ladder.
- PAPER_1166 — V(UA) Mexican-hat closure; $\langle R_4 \rangle$ value.
- PAPER_1167 — Master synthesis.
- PAPER_1170 — Four-line ledger; Route A.
- PAPER_1171 — KK regulator first-principles derivation.
- `uqff_closed_constants.py` — canonical integer-rational constants.