

stemflow: A Python Package for Adaptive Spatio-Temporal Exploratory Model

Yangkang Chen^{1,2}, Zhongru Gu^{1,3}, and Xiangjiang Zhan^{1,2,3,4}

¹ Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing, China ² University of Chinese Academy of Sciences, Beijing, China ³ Cardiff University-Institute of Zoology Joint Laboratory for Biocomplexity Research, Chinese Academy of Sciences, Beijing, China ⁴ Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming, China

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Stemflow is a user-friendly python package for Adaptive Spatio-Temporal Exploratory Model (AdaSTEM ([Fink et al., 2013](#))) that follows the style of scikit-learn BaseEstimator class ([Pedregosa et al., 2011](#)). It provides one-line model creation, fitting, prediction, and evaluation. It implements spatio-temporal train-test-split and cross-validation functions. After model training, feature importance could be evaluated with spatio-temporal dynamics. Stemflow also provides functions for visualizing ensembles structured in model training and generating GIF file for predicted results to animate the spatio-temporal movement of animal population.

Statement of need

Spatio-temporal big data is an emerging but valuable resource for ecological studies as human enter the era of big data ([Farley et al., 2018](#)). Data from broad-scale survey like citizen science projects is increasingly important in modern ecological research ([Dickinson et al., 2010](#)). Intensity of survey activities grows rapidly as more people are involved in citizen science in recent years, resulted in exponential accumulation of observational data ([Di Cecco et al., 2021](#); [Sullivan et al., 2014](#)). However, daily species observation records uploaded by non-professionals in citizen science program are known to have larger bias than professionally structured research, both in terms of data veracity and spatio-temporal balance of the datasets, which necessitates elaborate modeling methods to mine its merits ([Dickinson et al., 2010](#); [Farley et al., 2018](#)).

Some species distribution modeling (SDM) approaches were brought forward to adjust for bias in citizen science and model on the unobserved components ([Bird et al., 2014](#)). Still, many failed to account for the autocorrelation of space and time ([F. Dormann et al., 2007](#)), which is especially crucial in modeling inherently spatio-temporal biological events with variations at different scales ([Chave, 2013](#); [Levin, 1992](#)), such as seasonal migration. Adaptive Spatio-Temporal Exploratory Model (AdaSTEM) is a semi-parameterized machine learning model that leverages the spatio-temporal adjacency information of sample points to model occurrence or abundance of species ([Fink et al., 2013](#)). A QuadTree algorithm ([Samet, 1984](#)) is implemented to split data into smaller spatio-temporal grids (called stixels) conditional on the data abundance, with more abundant data allowing stixels to be divided into finer resolution (up to a maximum). Stixels with data size less than a certain threshold will not be modeled; instead, these stixels will be labeled as unpredictable. This procedure controls the degree of model extrapolation and reduces overfitting. A base model is trained for each stixel, that is, targets are only modeled on their adjacent information in space and time. Splitting-training is carried out several times to generate multiple ensembles. Finally, prediction results were aggregated across these ensembles.

43 AdaSTEM shows the capacity of supporting large scale spatio-temporal ecological data modeling
44 in many studies (Fink et al., 2020; Fuentes et al., 2023; La Sorte et al., 2022), especially for
45 modeling animal abundance at different scales (Fink et al., 2013). One well-known application
46 of AdaSTEM is the weekly abundance map of eBird Status and Trend product (Fink et al.,
47 2022), which was widely used as data sources of abundance data of bird populations (Bird
48 et al., 2014; Jarzyna & Stagge, 2023; Lin et al., 2022). The application of AdaSTEM could
49 be extended to other fields with similar data structure and spatio-temporal dependence, for
50 example, epidemiology. Despite the foreseeable significant role of spatio-temporal big data in
51 the coming decades of scientific research, the development of tools has not necessarily kept
52 pace.

53 Stemflow is positioned as a user-friendly python package to meet the need of general application
54 of modeling spatio-temporal large datasets. Scikit-learn style object-oriented modeling pipeline
55 enables concise model construction with compact parameterization at the user end, while the
56 rest of the modeling procedures are carried out under the hood. Once the fitting method is
57 called, the model class recursively splits the input training data into smaller spatio-temporal
58 stixels using QuadTree algorithm. For each of the stixels, a base model is trained only using
59 data falls into that stixel. Stixels are then aggregated and constitute an ensemble. In the
60 prediction phase, stemflow queries stixels for the input data according to their spatial and
61 temporal index, followed by corresponding base model prediction. Finally, prediction results
62 are aggregated across ensembles to generate robust estimations (see Fink et al. (2013) and
63 stemflow documentation for details).

64 For survey projects that include abundance information like eBird (Sullivan et al., 2014),
65 the targeted modeling values are often zero-inflated, owing to the fact of low observation
66 probability in many species. Zero-inflation could lead to poor regression model performance
67 (Campbell, 2021). In stemflow, we implement hurdle model classes that embed two sequential
68 models: a classifier to classify the absence and presence state, followed by a regressor to
69 model the abundance for prediction samples classified as presence. Hurdle model classes can
70 be conjunctively used with AdaSTEM model classes in two ways: Use hurdle model as the
71 base model for AdaSTEMRegressor (as in Johnston et al. (2015)), or use AdaSTEMClassifier
72 and AdaSTEMRegressor as the classifier and regressor in hurdle model. We demonstrate the
73 comparison of these two architectures in stemflow documentation.

74 One advantage of applying stemflow in scikit-learn style is that there is a variety of “base
75 models” to choose from scikit-learn or scikit-learn-style repertoire. The choices vary from
76 linear models to boosting and bagging tree-based models. Maxent model (C. B. Anderson,
77 2023) is also supported to play the role of “base model”, which largely expands the potential
78 application for presence-only modeling (see documentation).

79 While there exists mounting open source packages for species distribution modeling (mostly in
80 R, (Norberg et al., 2019); and one in Python (C. B. Anderson, 2023)), most of them solely
81 leverage environmental variables and do not support integration of spatio-temporal information
82 during model construction (but see C. B. Anderson (2023); S. C. Anderson et al. (2022);
83 Dobson et al. (2023)). This disadvantage is usually noted along with the overconfidence
84 of the model extrapolation capacity both for Maxent-based and ensemble-based models (A.
85 Lee-Yaw et al., 2022). To our knowledge, stemflow is the first package designed to solve the
86 spatio-temporal dependency conjugated with bias in data abundance distribution in species
87 distribution modeling. With the rapid accumulation of data and development of machine
88 learning techniques, stemflow will show potentials in spatio-temporal modeling, and could be
89 applied to other fields (e.g., epidemiology and weather prediction) in future.

90 Acknowledgements

91 This project was based upon work supported by National Natural Science Foundation of
92 China grants 32125005 (to X.Z.), 32270455 (to Z.G.), CAS Project for Young Scientists in

93 Basic Research (YSBR-097 to X.Z. and Z.G.). We thank Dr. Daniel Fink at Cornell Lab of
 94 Ornithology for the R scripts of STEM model and QuadTree which inspired our implementation
 95 in python.

96 References

- 97 A. Lee-Yaw, J., L. McCune, J., Pironon, S., & N. Sheth, S. (2022). Species distribution
 98 models rarely predict the biology of real populations. *Ecography*, 2022(6), e05877. <https://doi.org/10.1111/ecog.05877>
 99
- 100 Anderson, C. B. (2023). Elapid: Species distribution modeling tools for Python. *Journal of*
 101 *Open Source Software*, 8(84), 4930. <https://doi.org/10.21105/joss.04930>
- 102 Anderson, S. C., Ward, E. J., English, P. A., & Barnett, L. A. K. (2022). *sdmTMB: An R*
 103 *package for fast, flexible, and user-friendly generalized linear mixed effects models with*
 104 *spatial and spatiotemporal random fields* [Preprint]. *Ecology*. [https://doi.org/10.1101/](https://doi.org/10.1101/2022.03.24.485545)
 105 [2022.03.24.485545](https://doi.org/10.1101/2022.03.24.485545)
- 106 Bird, T. J., Bates, A. E., Lefcheck, J. S., Hill, N. A., Thomson, R. J., Edgar, G. J., Stuart-Smith,
 107 R. D., Wotherspoon, S., Krkosek, M., Stuart-Smith, J. F., Pecl, G. T., Barrett, N., &
 108 Frusher, S. (2014). Statistical solutions for error and bias in global citizen science datasets.
 109 *Biological Conservation*, 173, 144–154. <https://doi.org/10.1016/j.biocon.2013.07.037>
- 110 Campbell, H. (2021). The consequences of checking for zero-inflation and overdispersion in
 111 the analysis of count data. *Methods in Ecology and Evolution*, 12(4), 665–680. <https://doi.org/10.1111/2041-210X.13559>
 112
- 113 Chave, J. (2013). The problem of pattern and scale in ecology: What have we learned in 20
 114 years? *Ecology Letters*, 16(s1), 4–16. <https://doi.org/10.1111/ele.12048>
- 115 Di Cecco, G. J., Barve, V., Belitz, M. W., Stucky, B. J., Guralnick, R. P., & Hurlbert, A. H.
 116 (2021). Observing the Observers: How Participants Contribute Data to iNaturalist and
 117 Implications for Biodiversity Science. *BioScience*, 71(11), 1179–1188. [https://doi.org/10.](https://doi.org/10.1093/biosci/biab093)
 118 [1093/biosci/biab093](https://doi.org/10.1093/biosci/biab093)
- 119 Dickinson, J. L., Zuckerberg, B., & Bonter, D. N. (2010). Citizen Science as an Ecological
 120 Research Tool: Challenges and Benefits. *Annual Review of Ecology, Evolution, and*
 121 *Systematics*, 41(1), 149–172. <https://doi.org/10.1146/annurev-ecolsys-102209-144636>
- 122 Dobson, R., Challinor, A. J., Cheke, R. A., Jennings, S., Willis, S. G., & Dallimer, M. (2023).
 123 dynamicSDM: An r package for species geographical distribution and abundance modelling
 124 at high spatiotemporal resolution. *Methods in Ecology and Evolution*, 14(5), 1190–1199.
 125 <https://doi.org/10.1111/2041-210X.14101>
- 126 F. Dormann, C., M. McPherson, J., B. Araújo, M., Bivand, R., Bolliger, J., Carl, G., G. Davies,
 127 R., Hirzel, A., Jetz, W., Daniel Kissling, W., Kühn, I., Ohlemüller, R., R. Peres-Neto, P.,
 128 Reineking, B., Schröder, B., M. Schurr, F., & Wilson, R. (2007). Methods to account for
 129 spatial autocorrelation in the analysis of species distributional data: A review. *Ecography*,
 130 30(5), 609–628. <https://doi.org/10.1111/j.2007.0906-7590.05171.x>
- 131 Farley, S. S., Dawson, A., Goring, S. J., & Williams, J. W. (2018). Situating Ecology as
 132 a Big-Data Science: Current Advances, Challenges, and Solutions. *BioScience*, 68(8),
 133 563–576. <https://doi.org/10.1093/biosci/biy068>
- 134 Fink, D., Auer, T., Johnston, A., Ruiz-Gutierrez, V., Hochachka, W. M., & Kelling, S. (2020).
 135 Modeling avian full annual cycle distribution and population trends with citizen science
 136 data. *Ecological Applications*, 30(3), e02056. <https://doi.org/10.1002/eap.2056>
- 137 Fink, D., Auer, T., Johnston, A., Strimas-Mackey, M., Ligocki, S., Robinson, O., Hochachka,
 138 W., L. Jaromczyk, Rodewald, A., Wood, C., Davies, I., & Spencer., A. (2022). *eBird status*

- and trends, data version: 2021; released: 2022. Cornell Lab of Ornithology, Ithaca, New York. <https://doi.org/10.2173/ebirdst.2021>
- Fink, D., Damoulas, T., & Dave, J. (2013). Adaptive Spatio-Temporal Exploratory Models: Hemisphere-wide species distributions from massively crowdsourced eBird data. *Proceedings of the AAAI Conference on Artificial Intelligence*, 27(1), 1284–1290. <https://doi.org/10.1609/aaai.v27i1.8484>
- Fuentes, M., Van Doren, B. M., Fink, D., & Sheldon, D. (2023). BirdFlow: Learning seasonal bird movements from eBird data. *Methods in Ecology and Evolution*, 14(3), 923–938. <https://doi.org/10.1111/2041-210X.14052>
- Jarzyna, M. A., & Stagge, J. H. (2023). Decoupled spatiotemporal patterns of avian taxonomic and functional diversity. *Current Biology*, 33(6), 1153–1161.e4. <https://doi.org/10.1016/j.cub.2023.01.066>
- Johnston, A., Fink, D., Reynolds, M. D., Hochachka, W. M., Sullivan, B. L., Bruns, N. E., Hallstein, E., Merrifield, M. S., Matsumoto, S., & Kelling, S. (2015). Abundance models improve spatial and temporal prioritization of conservation resources. *Ecological Applications*, 25(7), 1749–1756. <https://doi.org/10.1890/14-1826.1>
- La Sorte, F. A., Horton, K. G., Johnston, A., Fink, D., & Auer, T. (2022). Seasonal associations with light pollution trends for nocturnally migrating bird populations. *Ecosphere*, 13(3), e3994. <https://doi.org/10.1002/ecs2.3994>
- Levin, S. A. (1992). The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award Lecture. *Ecology*, 73(6), 1943–1967. <https://doi.org/10.2307/1941447>
- Lin, H.-Y., Binley, A. D., Schuster, R., Rodewald, A. D., Buxton, R., & Bennett, J. R. (2022). Using community science data to help identify threatened species occurrences outside of known ranges. *Biological Conservation*, 268, 109523. <https://doi.org/10.1016/j.biocon.2022.109523>
- Norberg, A., Abrego, N., Blanchet, F. G., Adler, F. R., Anderson, B. J., Anttila, J., Araújo, M. B., Dallas, T., Dunson, D., Elith, J., Foster, S. D., Fox, R., Franklin, J., Godsoe, W., Guisan, A., O'Hara, B., Hill, N. A., Holt, R. D., Hui, F. K. C., ... Ovaskainen, O. (2019). A comprehensive evaluation of predictive performance of 33 species distribution models at species and community levels. *Ecological Monographs*, 89(3), e01370. <https://doi.org/10.1002/ecm.1370>
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., & Duchesnay, Édouard. (2011). Scikit-learn: Machine learning in python. *Journal of Machine Learning Research*, 12(85), 2825–2830. <http://jmlr.org/papers/v12/pedregosa11a.html>
- Samet, H. (1984). The Quadtree and Related Hierarchical Data Structures. *ACM Computing Surveys*, 16(2), 187–260. <https://doi.org/10.1145/356924.356930>
- Sullivan, B. L., Aycrigg, J. L., Barry, J. H., Bonney, R. E., Bruns, N., Cooper, C. B., Damoulas, T., Dhondt, A. A., Dietterich, T., Farnsworth, A., Fink, D., Fitzpatrick, J. W., Fredericks, T., Gerbracht, J., Gomes, C., Hochachka, W. M., Iliiff, M. J., Lagoze, C., La Sorte, F. A., ... Kelling, S. (2014). The eBird enterprise: An integrated approach to development and application of citizen science. *Biological Conservation*, 169, 31–40. <https://doi.org/10.1016/j.biocon.2013.11.003>